VOL. 667, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A APRIL 29, 1994

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Reproducibility of physical characteristics, protein immobilization and chromatographic performance of 3M Emphaze Biosupport Medium AB 1
by P.R. Johnson, N.J. Stern, P.D. Eitzman, J.K. Rasmussen, D.S. Milbrath, R.M. Gleason and R.E. Hogancamp (St. Paul, MN, USA) (Received November 4th, 1993) 1

6,6′-Dinitrobiphenyl-2,2′-dicarboxylic acid ionically bonded to aminopropyl silica: new axially chiral phase of C_2 symmetry for high-performance liquid chromatographic separation of enantiomeric amino alcohol derivatives
by M. Tichý, J. Holanová and J. Závada (Prague, Czech Republic) (Received November 9th, 1993) 11

Effect of mobile phase composition on the enantioselectivity of chromatographic separation on a quinine-bonded silica stationary phase
by P.N. Nesterenko, V.V. Krotov and S.M. Staroverov (Moscow, Russian Federation) (Received December 10th, 1993) 19

Determination of distribution coefficients for some 5-HT_3 receptor antagonists by reversed-phase high-performance liquid chromatography
by C. Kugel, B. Heintzelmann and J. Wagner (Strasbourg, France) (Received December 20th, 1993) 29

Hydrophobicity parameters of 2-chloro-2′-deoxyadenosine and some related analogues and retention in reversed-phase liquid chromatography
by V. Reichelová (Stockholm, Sweden and Bratislava, Slovak Republic) and F. Albertioni and J. Liliemark (Stockholm, Sweden) (Received November 23rd, 1993) 37

Ultra-trace analysis of phenols in water using high-performance liquid chromatography with on-line reaction detection
by G. Lamprecht and J.F.K. Huber (Vienna, Austria) (Received December 1st, 1993) 47

High-performance liquid chromatographic determination of naturally occurring flavonoids in *Citrus* with a photodiode-array detector
by Y. Nogata, H. Ohta and K.-I. Yoza (Hiroshima, Japan) and M. Berhow and S. Hasegawa (Pasadena, CA, USA) (Received December 22nd, 1993) 59

Separation of lactose, lactobionic acid and lactobionolactone by high-performance liquid chromatography
by P.J. Simms, K.B. Hicks, R.M. Haines, A.T. Hotchkiss, Jr. and S.F. Osman (Philadelphia, PA, USA) (Received December 20th, 1993) 67

Improved fractionation of sialylated glycopeptides by pellicular anion-exchange chromatography
by J.S. Rohrer (Sunnyvale, CA, USA) (Received December 24th, 1993) 75

Enantiomeric purity determination by high-performance liquid chromatography with coupled polarized photometric/UV detection. Analysis of adulterative addition of synthetic malic and tartaric acids
by A. Yamamoto, A. Matsunaga and E. Mizukami (Toyama, Japan) and K. Hayakawa and M. Miyazaki (Kanazawa, Japan) (Received November 17th, 1993) 85

Monitoring carboxylic acid formation in engine oils by liquid chromatography with fluorescence detection
by S.W. Lewis and P.J. Worsfold (Plymouth, UK) and E.H. McKerrell (Chester, UK) (Received December 24th, 1993) 91

Luminol chemiluminescent determination of hydrogen peroxide at picomole levels using high-performance liquid chromatography with a cation-exchange resin gel column
by T. Miyazawa, S. Lertsiri, K. Fujimoto and M. Oka (Sendai, Japan) (Received December 28th, 1993) 99

Determination of protein amino acids as butylthiocarbamyl derivatives by reversed-phase high-performance liquid chromatography with precolumn derivatization and UV detection
by K.-L. Woo and S.-H. Lee (Masan, South Korea) (Received October 26th, 1993) 105

(Continued overleaf)

Contents (continued)

Liquid chromatographic determination of carnitine by precolumn derivatization with pyrene-1-carbonyl cyanide
by K. Kamata, M. Takahashi, K. Terasima and M. Nishijima (Tokyo, Japan) (Received December 24th, 1993) . . 113

Effect of mobile phase composition on the separation of thyrotropin-releasing hormone and some metabolites by reversed-phase ion-pair chromatography
by S. Contreras Martinez (Cuernavaca Mor., Mexico) and L.E. Vera-Avila (Mexico, D.F., Mexico) (Received November 26th, 1993) 119

Rapid purification method for human recombinant tumor necrosis factor alpha
by A. Paquet, A. Lévesque and M. Pagé (Sainte-Foy, Canada) (Received December 31st, 1993) 125

Hydrophobic interaction chromatography for the purification of cytolytic bacterial toxins
by B. Schoel, M. Welzel and S.H.E. Kaufmann (Ulm, Germany) (Received November 30th, 1993) 131

New approaches for separating and purifying apple polyphenol oxidase isoenzymes: hydrophobic, metal chelate and affinity chromatography
by F. Richard-Forget and P. Goupy (Montfavet, France) and J. Nicolas (Paris, France) (Received December 16th, 1993) 141

Determination of N-methylcarbamate pesticides in environmental water samples using automated on-line trace enrichment with exchangeable cartridges and high-performance liquid chromatography
by M. Hiemstra and A. de Kok (Alkmaar, Netherlands) (Received January 6th, 1994) 155

Separation of acetylated polypropylene glycol di- and triamines by gradient reversed-phase high-performance liquid chromatography and evaporative light scattering detection
by K. Rissler (Basle, Switzerland) (Received January 11th, 1994) 167

Stability-indicating method for the determination of levodopa, levodopa–carbidopa and related impurities
by J.B. Kafil and B.S. Dhingra (Nutley, NJ, USA) (Received November 19th, 1993) 175

Direct optical resolution of vesamicol and a series of benzovesamicol analogues by high-performance liquid chromatography
by D.L. Gildersleeve, Y.-W. Jung and D.M. Wieland (Ann Arbor, MI, USA) (Received December 6th, 1993) . . 183

Gas Chromatography

Constancy of spectral response ratios in the flame photometric detector
by X.-Y. Sun and W.A. Aue (Halifax, Canada) (Received January 20th, 1994) 191

Comparison of universal chromatographic detectors for trace gas analysis
by R.T. Talasek and M.P. Schoenke (Dallas, TX, USA) (Received December 29th, 1993) 205

Design and construction of a simple supercritical fluid extraction system with semi-preparative and preparative capabilities for application to natural products
by S.R. Sargenti and F.M. Lanças (São Carlos, Brazil) (Received December 20th, 1993) 213

Determination of sulphonic compounds as their thiotrifluoroacetate derivatives by gas chromatography with ion trap detection
by X.-d. Li and Z.-s. Lin (Shanghai, China) (Received November 23rd, 1993) 219

Determination of neutral lipids from subcutaneous fat of cured ham by capillary gas chromatography and liquid chromatography
by J.A. García Regueiro, J. Gibert and I. Díaz (Girona, Spain) (Received December 6th, 1993) 225

Residual solvent analysis by headspace gas chromatography
by N. Kumar and J.G. Gow (Collegeville, PA, USA) (Received December 13th, 1993) 235

Determination of organophosphorus and triazine pesticides in ground- and drinking water by solid-phase extraction and gas chromatography with nitrogen–phosphorus or mass spectrometric detection
by M. Psathaki, E. Manoussaridou and E.G. Stephanou (Heraclion, Greece) (Received October 12th, 1993) . . . 241

Supercritical Fluid Chromatography

Effect of temperature and density on the performance of micropacked columns in supercritical fluid chromatography
by E. Ibáñez, J. Tabera, M. Herraiz and G. Reglero (Madrid, Spain) (Received October 19th, 1993) 249

Electrophoresis

Computer simulation for capillary zone electrophoresis. A quantitative approach
by S.V. Ermakov and P.G. Righetti (Milan, Italy) (Received November 15th, 1993) 257

Capillary zone electrophoresis of synthetic opioid and tachykinin peptides
by H.G. Lee and D.M. Desiderio (Memphis, TN, USA) (Received November 11th, 1993) 271

(Continued overleaf)

SHORT COMMUNICATIONS

Column Liquid Chromatography

Reversed-phase high-performance liquid chromatography of functionalized dendritic macromolecules
by J.M.J. Fréchet, L. Lochman, V. Šmigol and F. Švec (Ithaca, NY, USA) (Received November 24th, 1993) . . . 284

Identification of flavan-3-ols and procyanidins by high-performance liquid chromatography and chemical reaction detection
by D. Treutter, C. Santos-Buelga and M. Gutmann (Freising-Weihenstephan, Germany) and H. Kolodziej (Berlin, Germany) (Received January 13th, 1994) . . . 290

Simultaneous analysis of hydroxylamine, N-methylhydroxylamine and N,N-dimethylhydroxylamine by ion chromatography
by A.M. Prokai and R.K. Ravichandran (Indianapolis, IN, USA) (Received January 28th, 1994) . . . 298

Peptide mapping using combinations of size-exclusion chromatography, reversed-phase chromatography and capillary electrophoresis
by M. Strömqvist (Umeå, Sweden) (Received February 14th, 1994) . . . 304

Chromatographic evaluation of the binding of lysozyme to poly(dimethyldiallylammonium chloride)
by J. Xia and P.L. Dubin (Indianapolis, IN, USA) (Received January 5th, 1994) . . . 311

Comparison of extraction procedures for high-performance liquid chromatographic determination of cellular deoxynucleotides
by S. Palmer (Huddinge, Sweden) and S. Cox (Huddinge and Stockholm, Sweden) (Received January 18th, 1994) 316

Application of centrifugal partition chromatography to the separation of Lauraceous alkaloids
by S.-S. Lee (Taipei, Taiwan) (Received January 17th, 1994) . . . 322

Application of high-performance liquid chromatography to the determination of bitter principles of pharmaceutical relevance
by O.P. Semenova, A.R. Timerbaev and G.K. Bonn (Linz, Austria) (Received November 23rd, 1993) . . . 327

Gas Chromatography

Application of a fused-silica column to the determination of very volatile amines by gas–solid chromatography
by M. Mohnke, B. Schmidt and R. Schmidt (Leipzig, Germany) and J.C. Buijten and Ph. Mussche (Middelburg, Netherlands) (Received January 31st, 1994) . . . 334

Influence of dissolved gases in the dynamic headspace analysis of styrene and other volatile organic compounds and improvement of their determination
by J. Le Sech, V. Ducruet and A. Feigenbaum (Jouy en Josas, France) (Received December 27th, 1993) . . . 340

Rapid method for the determination of multiple pyrethroid residues in fruits and vegetables by capillary column gas chromatography
by G.-F. Pang, C.-L. Fan, Y.-Z. Chao and T.-S. Zhao (Hebei, China) (Received January 17th, 1994) . . . 348

Supercritical Fluid Chromatography

Molar mass determination of oligomeric ethylene oxide adducts using supercritical fluid chromatography and matrix-assisted laser desorption–ionization time-of-flight mass spectrometry
by U. Just and H.-R. Holzbauer (Berlin, Germany) and M. Resch (Duisburg, Germany) (Received November 15th, 1993) . . . 354

Planar Chromatography

Thin-layer chromatographic determination of β-carotene, cantaxanthin, lutein, violaxanthin and neoxanthin on Chromarods
by A.J. Rosas Romero, J.C. Herrera, E. Martinez De Aparicio and E.A. Molina Cuevas (Caracas, Venezuela) (Received November 29th, 1993) . . . 361

Electrophoresis

Separation of chiral isomers of *p*-nitrophenyl-2-amino-3-hydroxypropanone by capillary zone electrophoresis using cyclodextrins as chiral selector
by J. Gu and R. Fu (Beijing, China) (Received December 7th, 1993) . . . 367

Ionophoretic technique in the study of Cu(II), Ni(II), Co(II) and Mn(II)–hydrazine–nitrilotriacetate mixed-ligand complexes
by U. Mishra and R.K.P. Singh (Allahabad, India) (Received January 17th, 1994) . . . 371

AUTHOR INDEX . . . 377